

# DuPont™ Vydate® L

insecticide/nematicide



## Safe Use and Handling of Vydate® L



1. Keep pesticides out of reach of children and animals in a locked, preferably bunded storage area.



2. Before using Vydate® L read the label carefully.



3. DO NOT use the application equipment if it is leaky or damaged. Always check to ensure that the application equipment is in proper working condition.



4. Avoid contact with Vydate® L at all times.



5. Always wash hands, arms and face thoroughly with soap and water before eating, drinking, smoking, chewing gum or using the toilet.



6. If you feel sick stop applying and seek urgent medical attention.



7. Triple rinse empty non-returnable containers and send them to an accredited recycling site. Dispose of rinsate in the field or in a designated safe disposal pit.



8. DO NOT reuse empty Vydate® L containers.

## Correct Protective Gear

Elbow length PVC gloves

Cotton overalls buttoned to neck and wrist

Impervious footwear




## Symptoms of Poisoning

Poisoning produces effects associated with anticholinesterase activity which may include:


- Weakness
- Blurred vision
- Breathing difficulties
- Nausea
- Headache
- Abdominal pain
- Discomfort in the chest
- Constriction of pupils
- Slow pulse
- Sweating
- Muscle twitching

## First Aid for Poisoning


If poisoning occurs, contact a doctor or the Poisons Information Centre on 13 11 26. Have the product container or label with you when calling a doctor, Poisons Information Centre or going for treatment.



EYE CONTACT




Hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.



SWALLOWED



If swallowed, activated charcoal may be advised. Give atropine if instructed.



SKIN CONTACT



Take off all contaminated clothing immediately. In case of skin contact, wash off immediately with plenty of water.



INHALATION



Remove person to fresh air. If signs/symptoms continue, get medical attention. Artificial respiration and/or oxygen may be necessary.

IN AN  
EMERGENCY  
CALL

000

